

Quality ID #128: Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan

2026 COLLECTION TYPE:

MERIT-BASED INCENTIVE PAYMENT SYSTEM (MIPS) CLINICAL QUALITY MEASURE (CQM)
– MIPS VALUE PATHWAY (MVP) REPORTING ONLY

MEASURE TYPE:

Process

DESCRIPTION:

Percentage of patients aged 18 years and older with a BMI documented during the current encounter or within the previous twelve months AND who had a follow-up plan documented if the most recent BMI was outside of normal parameters.

INSTRUCTIONS:

Reporting Frequency:

This measure is to be submitted a minimum of once per performance period for denominator eligible cases as defined in the denominator criteria.

Intent and Clinician Applicability:

This measure is intended to reflect the quality of services provided for patients seen during the performance period. This measure may be submitted by Merit-based Incentive Payment System (MIPS) eligible clinicians who perform the quality actions as defined by the numerator based on the services provided and the measure-specific denominator coding.

Measure Strata and Performance Rates:

This measure contains one strata defined by a single submission criteria.

This measure produces a single performance rate.

Implementation Considerations:

For the purposes of MIPS implementation, this patient-intermediate measure is submitted a minimum of once per patient for the performance period. The most recent numerator option/quality data code will be used if the measure is submitted more than once.

There is no diagnosis associated with this measure. The BMI may be documented in the medical record of the provider or in outside medical records obtained by the provider. If the most recent documented BMI is outside of normal parameters, then a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter. The documented follow-up plan must be based on the most recent documented BMI outside of normal parameters, example: "Patient referred to nutrition counseling for BMI above or below normal parameters" (See Definitions for examples of follow-up plan treatments). If more than one BMI is submitted during the measurement period, the most recent BMI will be used to determine if the performance has been met. Review the exclusions and exceptions criteria to determine those patients that BMI measurement may not be appropriate or necessary.

This measure specification is only available for MIPS Value Pathway (MVP) reporting and is not available for traditional MIPS reporting.

Telehealth:

NOT TELEHEALTH ELIGIBLE: This measure is not appropriate for nor applicable to the telehealth setting.

Patient encounters for this measure conducted via telehealth should be removed from the denominator eligible patient population. Therefore, if the patient meets all denominator criteria but the encounter is conducted via telehealth, it would be appropriate to remove them from the denominator eligible patient population. Telehealth eligibility is at the measure level for inclusion within the denominator eligible patient population and based on the measure specification definitions which are independent of changes to coding and/or billing practices.

Measure Submission:

The quality data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third party intermediaries that utilize this collection type for submissions; however, these codes may be submitted for those third party intermediaries that utilize Medicare Part B claims data. The coding provided to identify the measure criteria: Denominator or Numerator, may be an example of coding that could be used to identify patients that meet the intent of this clinical topic. When implementing this measure, please refer to the 'Reference Coding' section to determine if other codes or code languages that meet the intent of the criteria may also be used within the medical record to identify and/or assess patients. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

DENOMINATOR:

All patients aged 18 and older on the date of the encounter with at least one qualifying encounter during the measurement period.

Definition:

Not Eligible for BMI Screening or Follow-Up Plan (Denominator Exclusions) – A patient is not eligible if one or more of the following reasons are documented:

- Patients receiving palliative or hospice care on the date of the current encounter or any time prior to the current encounter
- Patients who are pregnant on the date of the current encounter or any time during the measurement period prior to the current encounter

DENOMINATOR NOTE:

**Signifies that this CPT Category I code is a non-covered service under the Medicare Part B Physician Fee Schedule (PFS). These non-covered services should be counted in the denominator population for MIPS CQMs.*

Denominator Criteria (Eligible Cases):

Patients aged \geq 18 years on date of encounter

AND

Patient encounter during the performance period (CPT or HCPCS): 90791, 90792, 90832, 90834, 90837, 96156, 96158, 97161, 97162, 97163, 97165, 97166, 97167, 97802, 97803, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99304, 99305, 99306, 99307, 99308, 99309, 99310, 99315, 99316, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99385*, 99386*, 99387*, 99395*, 99396*, 99397*, 99401*, 99402*, 99424, 99491, D7111, D7140, D7210, D7220, D7230, D7240, D7241, D7250, D7251, G0101, G0108, G0270, G0271, G0402, G0438, G0439, G0447, G0473

WITHOUT

Encounters conducted via telehealth: M1431

WITHOUT

Place of Service (POS): 12

AND NOT

DENOMINATOR EXCLUSIONS:

Documentation stating the patient has received or is currently receiving palliative or hospice care: G9996

OR

Documentation of patient pregnancy anytime during the measurement period prior to and including the current encounter: G9997

NUMERATOR:

Patients with a documented BMI during the encounter or during the previous twelve months AND when the BMI is outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the encounter.

Definitions:

Normal BMI Parameters – Age 18 years and older $BMI \geq 18.5$ and $< 25 \text{ kg/m}^2$

BMI – Body mass index (BMI) is a number calculated using the Quetelet index: weight divided by height squared (W/H^2) and is commonly used to classify weight categories. "BMI" can be calculated using:

Metric Units: $BMI = \text{Weight (kg)} / (\text{Height (m)} \times \text{Height (m)})$

OR

English Units: $BMI = \text{Weight (lbs)} / (\text{Height (in)} \times \text{Height (in)}) \times 703$

Follow-Up Plan – Proposed outline of treatment to be conducted as a result of a BMI outside of normal parameters. A "follow-up" plan may include, but is not limited to:

- Documentation of education
- Referral (for example a Registered Dietitian Nutritionist (RDN), occupational therapist, physical therapist, primary care provider, exercise physiologist, mental health professional, or surgeon) for lifestyle/behavioral therapy
- Pharmacological interventions
- Dietary supplements
- Exercise counseling
- Nutrition counseling

Patients with a Documented Reason for Not Screening BMI (Denominator Exception) -

Patient Reason:

- Patients who refuse measurement of height and/or weight on the date of the current encounter or any time during the measurement period prior to the current encounter

OR

Medical Reason:

- Patients with a documented medical reason for not documenting BMI such as patients in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient's health status.

Patients with a Documented Reason for Not Documenting a Follow-up Plan for BMI Outside Normal Parameters (Denominator Exception) - Medical Reason(s):

- Patients (e.g., elderly patients 65 years of age or older) for whom weight reduction/weight gain would complicate other underlying health conditions such as illness or physical disability, mental illness, dementia, confusion, or nutritional deficiency such as vitamin/mineral deficiency; patients in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient's health status

Numerator Instructions:

Height and Weight - An eligible clinician or their staff is required to measure both height and weight. Both height and weight must be measured within twelve months of the current encounter. Self-reported values cannot be used.

- The BMI may be documented in the medical record of the provider or in outside medical records obtained by the provider.
- If more than one BMI is reported during the measurement period, the most recent BMI will be used to determine if the performance has been met.

Follow-Up Plan – If the most recent documented BMI is outside of normal parameters, then a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter. The documented follow-up plan must be based on the most recent documented BMI, outside of normal parameters, example: “Patient referred to nutrition counseling for BMI above or below normal parameters”. (See Definitions for examples of follow-up plan treatments).

Performance Met for G8417 & G8418 –

- If the provider documents a BMI and a follow-up plan for a BMI outside normal parameters at the current encounter **OR**
- If the patient has a documented BMI within the previous twelve months of the current encounter, the provider documents a follow-up plan for a BMI outside normal parameters at the current encounter **OR**
- If the patient has a documented BMI within the previous twelve months of the current encounter **AND** the patient has a documented follow-up plan for a BMI outside normal parameters within the previous twelve months of the current encounter

Numerator Options:

Performance Met:

BMI is documented within normal parameters and no follow-up plan is required (G8420)

OR

Performance Met:

BMI is documented above normal parameters and a follow-up plan is documented (G8417)

OR

Performance Met:

BMI is documented below normal parameters and a follow-up plan is documented (G8418)

OR

Denominator Exception:

BMI not documented due to medical reason OR patient refusal of height or weight measurement (G2181)

OR

Denominator Exception:

BMI is documented as being outside of normal parameters, follow-up plan is not completed for documented medical reason (G9716)

OR

Performance Not Met:

BMI not documented and no reason is given (G8421)

OR

Performance Not Met:

BMI documented outside normal parameters, no follow-up plan documented, no reason given (G8419)

RATIONALE:

BMI Above Normal Parameters

“Obesity is a chronic, multifactorial disease with complex psychological, environmental (social and cultural), genetic, physiologic, metabolic and behavioral causes and consequences. The prevalence of overweight and obese people is increasing worldwide at an alarming rate in both developing and developed countries. Environmental and behavioral changes brought about by economic development, modernization and urbanization have been linked to the rise in global obesity. The health consequences are becoming apparent (1).”

More than a third of U.S. adults have a body mass index (BMI) $\geq 30 \text{ kg/m}^2$ and are at increased risk for diabetes, cardiovascular disease (CVD), and obstructive sleep apnea (2,3, 4). Hales reported that the prevalence of obesity among adults and youth in the United States was 39.8 percent and 18.5 percent respectively, from 2015–2016. Furthermore, the prevalence of obesity in adults increased to 42.4 percent in 2018, with the highest percentage among adults in the 40–59 age bracket compared with other age groups (5). Hales also disaggregated the data according to race/ethnicity and noted that obesity prevalence was higher among non-Hispanic Black adults and Hispanic adults when compared with other races and ethnicities. Obesity prevalence was lowest among non-Hispanic Asian men and women. Among men, obesity prevalence was higher among Hispanic men compared with non-Hispanic Black men and non-Hispanic White men. Obesity prevalence was higher among Hispanic men compared with non-Hispanic Black men. Among women, the prevalence among non-Hispanic Black women was 56.9 percent, which was higher than all other race/ethnicities. In general, the prevalence of obesity in the U.S. remains higher than the Healthy People 2020 goal of 30.5 percent among adults (6).

BMI continues to be a common and reasonably reliable measurement to identify overweight and obese adults who may be at an increased risk for future morbidity. Although good quality evidence supports obtaining a BMI, it is important to recognize it is not a perfect measurement. For example, BMI and its associated disease and mortality risk appear to vary among ethnic subgroups. Black/African Americans appear to have the lowest mortality risk at a BMI of 26.2-28.5 kg/m² in Black women and 27.1-30.2 kg/m² in Black men. In contrast, Asian populations may experience lowest mortality rates starting at a BMI of 23 to 24 kg/m². The correlation between BMI and diabetes risk also varies by ethnicity (7). Moreover, BMI is not a direct measure of adiposity and as a consequence, it can over or underestimate adiposity. However, overall, BMI is a derived value that correlates well with total body fat and markers of secondary complications, e.g., hypertension and dyslipidemia (8).

Furthermore, it is important to enhance beneficiary access to appropriate treatments for obesity, which could result in decreased healthcare costs and lower obesity rates. Behavioral weight management treatment has been identified as an effective first-line treatment for obesity with an average initial weight loss of 8-10 percent. This percentage of weight loss is associated with a significant risk reduction for diabetes and CVD (9). Evidence also shows that when provided 14 or more high-intensity behavioral intervention sessions of face-to-face individual or group treatment across 6 months, participants lose up to 8 percent of their weight during that time and experience improvements in heart disease risk factors and quality of life (10). There is also evidence that high-intensity behavioral counseling is effective, whether delivered in-person, by phone, or electronically (11). Moreover, Intensive Behavioral Therapy (IBT) for obesity provided by Registered Dietitian Nutritionists for 6-12 months shows significant mean weight loss of up to 10 percent of body weight, maintained over one year's time (12).

Despite the evidence that supports weight management counseling, the rate of use in primary care for patients with obesity decreased by 10 percent from 39.9 percent in 1995-1996 to 29.9 percent in 2007-2008 (13). Weight management counseling during primary care visits further declined from 33 percent to 21 percent between 2008-2009 and 2012-2013. This suggests that obesity management in primary care remains suboptimal (14). Therefore, screening for BMI and follow-up is critical and will help in reaching the quality goals of population health and cost reduction. However, due to concerns for other underlying conditions (such as bone health) or nutrition related deficiencies providers are cautioned to use their best clinical judgment and when considering weight management programs for overweight patients, especially the elderly (15).

BMI Below Normal Parameters

On the other end of the body weight spectrum is underweight (BMI $< 18.5 \text{ kg/m}^2$), which is also detrimental to population health. When compared to normal weight individuals (BMI 18.5-25 kg/m²), underweight individuals have significantly higher death rates with a Hazard Ratio of 2.27 and 95 percent confidence intervals (CI) = 1.78, 2.90 (16). Individuals with a BMI $< 18.5 \text{ kg/m}^2$ have been shown to be at a higher risk for adverse events, postoperative infection, and/or mortality following a surgical procedure (17, 18, 19, 20). BMI below normal parameters is a risk factor for developing severe illness from respiratory infections such as influenza and COVID-19 (21, 22). BMI below normal parameters can negatively impact both male and female fertility (23, 24).

Poor nutrition or underlying health conditions can result in underweight (25). The National Health and Nutrition Examination Survey (NHANES) results from 2007-2010 indicate that women are more likely to be underweight than men (25). However, all patients should be equally screened for underweight and followed up with nutritional counseling or another clinically appropriate intervention to reduce mortality and morbidity associated with underweight.

CLINICAL RECOMMENDATION STATEMENTS:

All adults should be screened annually using a BMI measurement. BMI measurements $\geq 25\text{kg/m}^2$ should be used to initiate further evaluation of overweight or obesity after taking into account age, gender, ethnicity, fluid status, and muscularity; therefore, clinical evaluation and judgment must be used when BMI is employed as the anthropometric indicator of excess adiposity, particularly in athletes and those with sarcopenia (26) (Grade A).

Overweight and Underweight Categories:

Underweight < 18.5 ; Normal weight 18.5-24.9; Overweight 25-29.9; Obese class I 30-34.9; Obese class II 35-39.9; Obese class III ≥ 40 (26).

BMI cutoff point value of $\geq 23\text{ kg/m}^2$ should be used in the screening and confirmation of excess adiposity in Asian adults (26) (Grade B).

Lifestyle/Behavioral Therapy for Overweight and Obesity should include behavioral interventions that enhance adherence to prescriptions for a reduced-calorie meal plan and increased physical activity (behavioral interventions can include: self-monitoring of weight, food intake, and physical activity; clear and reasonable goal-setting; education pertaining to obesity, nutrition, and physical activity; face-to-face and group meetings; stimulus control; systematic approaches for problem solving; stress reduction; cognitive restructuring [i.e., cognitive behavioral therapy], motivational interviewing; behavioral contracting; psychological counseling; and mobilization of social support structures) (26) (Grade A).

Behavioral lifestyle intervention should be tailored to a patient's ethnic, cultural, socioeconomic, and educational background (26) (Grade B).

The U.S. Preventive Services Task Force (USPSTF) recommends that clinicians offer or refer adults with a BMI of 30 kg/m^2 or higher to intensive, multicomponent behavioral interventions.

Interventions:

- Effective intensive behavioral interventions were designed to help participants achieve or maintain a weight loss of at least five percent through a combination of dietary changes and increased physical activity
- Most interventions lasted for one to two years, and the majority had at least 12 sessions in the first year
- Most behavioral interventions focused on problem solving to identify barriers, self-monitoring of weight, peer support, and relapse prevention
- Interventions also provided tools to support weight loss or weight loss maintenance (e.g., pedometers, food scales, or exercise videos) (Grade B) (27)

Nutritional safety for the elderly should be considered when recommending weight reduction. "A clinical decision to forego obesity treatment in older adults should be guided by an evaluation of the potential benefits of weight reduction for day-to-day functioning and reduction of the risk of future cardiovascular events, as well as the patient's motivation for weight reduction. Care must be taken to ensure that any weight reduction program minimizes the likelihood of adverse effects on bone health or other aspects of nutritional status" (15) (Evidence Category D). In addition, weight reduction prescriptions in older persons should be accompanied by proper nutritional counseling and regular body weight monitoring (15).

The possibility that a standard approach to weight loss will work differently in diverse patient populations must be considered when setting expectations about treatment outcomes (15) (Evidence Category B).

REFERENCES:

1. Fitch, A., Everling, L., Fox, C., Goldberg, J., Heim, C., Johnson, K., Kaufman, T., Kennedy, E., Kestenbaum, C., Lano, M., Leslie, D., Newell, T., O'Connor, P., Slusarek, B., Spaniol, A., Stovitz, S., & Webb, B. (2013). Prevention and management of obesity for adults. Bloomington, MN: Institute for Clinical Systems Improvement.
2. Flegal, K. M., Carroll, M. D., Kit, B. K., & Ogden, C. L. (2012). Prevalence of obesity and trends in the distribution of body mass index among U. S. adults, 1999-2010. *JAMA*, 307(5), 491-497. [doi:10.1001/jama.2012.39](https://doi.org/10.1001/jama.2012.39)
3. Ogden, C.L., Carroll, M.D., Fryar, C.D., Flegal, K.M. (2015). Prevalence of obesity among adults and youth: United States, 2011–2014. NCHS data brief, no 219. Hyattsville, MD: National Center for Health Statistics. Retrieved from <https://www.cdc.gov/nchs/data/databriefs/db219.pdf>
4. Dong, Z., Xu, X., Wang, C., Cartledge, S., Maddison, R., & Mohammed Shariful Islam, S. (2020). Association of overweight and obesity with obstructive sleep apnoea: A systematic review and meta-analysis. *Obesity Medicine*, 17. doi: <https://doi.org/10.1016/j.obmed.2020.100185>
5. Hales, C. M., Carroll, M. D., Fryar, C. D., et al. (2017). Prevalence of obesity among adults and youth: United States, 2015-2016. NCHS Data Brief No. 288. <https://www.cdc.gov/nchs/products/databriefs/db288.htm>
6. Hales, C. M., Carroll, M. D., Fryar, C. D., & Ogden, C.L. (2020). Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017-2018. NCHS Data Brief No. 360. <https://www.cdc.gov/nchs/products/databriefs/db360.htm>
7. LeBlanc, E., O'Connor, E., Whitlock, E. P., Patnode, C., & Kapka, T. (2011). Screening for and management of obesity and overweight in adults (Evidence Report No. 89; AHRQ Publication No. 11-05159-EF-1). Rockville, MD: Agency for Healthcare Research and Quality
8. Barlow, S. E., & the Expert Committee. (2007). Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. *Pediatrics*, 120(Suppl. 4), S164-S192. [doi:10.1542/peds.2007-2329C](https://doi.org/10.1542/peds.2007-2329C)
9. Wadden, T. A., Butrym, M. L., & Wilson, C. (2007). Lifestyle modification for the management of obesity. *Gastroenterology*, 132 (6), 2226-2238. doi: [10.1053/j.gastro.2007.03.051](https://doi.org/10.1053/j.gastro.2007.03.051)
10. Wadden, T. A., Tronieri, J. S., & Butrym, M. L. (2020). Lifestyle modification approaches for the treatment of obesity in adults. *American Psychologist*, 75(2), 235-251
11. Tronieri, J. S., Wadden, T. A., Chao, A. M., & Tsai, A. G. (2019). Primary Care Interventions for Obesity: Review of the Evidence. *Current obesity reports*, 8(2), 128–136. <https://doi.org/10.1007/s13679-019-00341-5>
12. Raynor, H. A., & Champagne, C. M. (2016). Position of the Academy of Nutrition and Dietetics: Interventions for the treatment of overweight and obesity in adults. *Journal of the Academy of Nutrition and Dietetics*, 116(1), 129-147. [doi:10.1016/j.jand.2015.10.031](https://doi.org/10.1016/j.jand.2015.10.031)
13. Kraschnewski, J. L., Sciamanna, C. N., Stuckey, H. L., Chuang, C. H., Lehman, E. B., Hwang, K. O., Sherwood, L. L., & Nembhard, H. B. (2013). A silent response to the obesity epidemic: decline in US physician weight counseling. *Medical care*, 51(2), 186–192. <https://doi.org/10.1097/MLR.0b013e3182726c33>
14. Fitzpatrick, S. L., & Stevens, V. J. (2017). Adult obesity management in primary care, 2008-2013. *Preventive medicine*, 99, 128-133. <https://doi.org/10.1016/j.ypmed.2017.02.020>
15. NHLBI Obesity Education Initiative. (1998). Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults (Report No. 98-4083). Bethesda, MD: NHLBI
16. Borrell, L. N., & Samuel, L. (2014). Body mass index categories and mortality risk in U.S. adults: The effect of overweight and obesity on advancing death. *American Journal of Public Health*, 104(3), 512-519. [doi:10.2105/AJPH.2013.301597](https://doi.org/10.2105/AJPH.2013.301597)
17. Katakam, A., Melnic, C. M., Bragdon, C. R., Sauder, N., Collins, A. K., & Bedair, H. S. (2021). Low body mass index is a predictor for mortality and increased length of stay following total joint arthroplasty. *The Journal of Arthroplasty*, 36(1), 72-77. <https://doi.org/10.1016/j.arth.2020.07.055>
18. Ottesen, T. D., Malpani, R., Galivanche, A. R., Zogg, C. K., Varthi, A. G., & Grauer, J. N. (2020). Underweight patients are at just as much risk as super morbidly obese patients when undergoing anterior cervical spine surgery. *The Spine Journal: Official Journal of the North American Spine Society*, 20(7), 1085-1095. <https://doi.org/10.1016/j.spinee.2020.03.007>

19. Ottesen, T. D., Galivanche, A. R., Greene, J. D., Malpani, R., Varthi, A. G., & Grauer, J. N. (2022). Underweight patients are the highest risk body mass index group for perioperative adverse events following stand-alone anterior lumbar interbody fusion. *The Spine Journal: Official Journal of the North American Spine Society*, 22(7), 1139-1148. <https://doi.org/10.1016/j.spinee.2022.02.012>
20. Rudasill, S. E., Dillon, D., Karunungan, K., Mardock, A. L., Hadaya, J., Sanailha, Y., Tran, Z., & Benharash, P. (2021). The obesity paradox: Underweight patients are at the greatest risk of mortality after cholecystectomy. *Surgery*, 170(3), 675-681. <https://doi.org/10.1016/j.surg.2021.03.034>
21. Moser, J. S., Galindo-Fraga, A., Ortiz-Hernández, A. A., Gu, W., Hunsberger, S., Galán-Herrera, J. F., Guerrero, M. L., Ruiz-Palacios, G. M., Beigel, J. H., & La Red ILI 002 Study Group. (2019). Underweight, overweight, and obesity as independent risk factors for hospitalization in adults and children from influenza and other respiratory viruses. *Influenza and Other Respiratory Viruses*, 13(1), 3-9. <https://doi.org/10.1111/irv.12618>
22. Ye, P., Pang, R., Li, L., Li, H. R., Liu, S. L., & Zhao, L. (2021). Both underweight and obesity are associated with an increased risk of coronavirus disease 2019 (COVID-19) severity. *Frontiers in Nutrition*, 8, 649422. <https://doi.org/10.3389/fnut.2021.649422>
23. Boutari, C., Pappas, P. D., Mintziori, G., Nigdelis, M. P., Athanasiadis, L., Goulis, D. G., Mantzoros, C. S. (2020). The effect of underweight on female and male reproduction. *Metabolism*, 107, 154229. <https://doi.org/10.1016/j.metabol.2020.154229>
24. Guo, D., Xu, M., Zhou, Q., Wu, C., Ju, R., & Dai, J. (2019). Is low body mass index a risk factor for semen quality? A PRISMA-compliant meta-analysis. *Medicine*, 98(32), e16677. <https://doi.org/10.1097/MD.00000000000016677>
25. Fryar, C. D., & Ogden, C. L. (2012). Prevalence of underweight among adults aged 20 and over: United States, 1960-1962 through 2007-2010. Hyattsville, MD: NCHS, Division of Health and Nutrition Examination Surveys. http://www.cdc.gov/nchs/data/hestat/underweight_adult_07_10/underweight_adult_07_10.pdf
26. Garvey, W. T., Mechanick, J. I., Brett, E. M., Garber, A. J., Hurley, D. L., Jastreboff, A. M., Nadolsky, K., Pessah-Pollack, R., Plodkowski, R., & Reviewers of the AACE/ACE Obesity Clinical Practice Guidelines. (2016). American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocrine Practice*, 22(Suppl. 3), 1-203. <doi:10.4158/EP161365GL>
27. U.S. Preventive Services Task Force (USPSTF). (2018). Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults: U.S. Preventive Services Task Force recommendation statement. *JAMA*, 320(11), 1163-1171. <doi:10.1001/jama.2018.13022>

COPYRIGHT:

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

©

This measure is owned and stewarded by the Centers for Medicare & Medicaid Services (CMS). CMS contracted (Contract # 75FCMC18D0027/ Task Order # 75FCMC24F0144) with the American Institutes for Research (AIR) to develop this measure. AIR is not responsible for any use of the Measure. AIR makes no representations, warranties, or endorsement about the quality of any organization or physician that uses or reports performance measures and AIR has no liability to anyone who relies on such measures or specifications. This measure is in the public domain.


Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. AIR disclaims all liability for use or accuracy of any third party codes contained in the specifications.


CPT® contained in the Measure specifications is copyright 2004-2025 American Medical Association. ICD-10 is copyright 2025 World Health Organization. All Rights Reserved.

These performance measures are not clinical guidelines and do not establish a standard of medical care, and have not been tested for all potential applications.

**2026 Clinical Quality Measure Flow for Quality ID #128:
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan**

Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.





SAMPLE CALCULATIONS

Data Completeness=

$$\frac{\text{Performance Met } (a^1 + a^2 + a^3 = 50 \text{ patients}) + \text{Denominator Exception } (b^1 + b^2 = 0 \text{ patients}) + \text{Performance Not Met } (c^1 + c^2 = 20 \text{ patients})}{\text{Eligible Population / Denominator } (d = 80 \text{ patients})} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

Performance Rate=

$$\frac{\text{Performance Met } (a^1 + a^2 + a^3 = 50 \text{ patients})}{\text{Data Completeness Numerator } (70 \text{ patients}) - \text{Denominator Exception } (b^1 + b^2 = 0 \text{ patients})} = \frac{50 \text{ patients}}{70 \text{ patients}} = 71.43\%$$

* See the posted measure specification for specific coding and instructions to submit this measure.
 NOTE: Submission Frequency: Patient-Intermediate

CPT only copyright 2025 American Medical Association. All rights reserved.
 The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

v10

2026 Clinical Quality Measure Flow Narrative for Quality ID #128:
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan

Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.

1. Start with Denominator
2. Check *Patients aged greater than or equal to 18 years on date of encounter*:
 - a. If *Patients aged greater than or equal to 18 years on date of encounter* equals No, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Patients aged greater than or equal to 18 years on date of encounter* equals Yes, proceed to check *Patient encounter during the performance period as listed in Denominator**.
3. Check *Patient encounter during the performance period as listed in Denominator**:
 - a. If *Patient encounter during the performance period as listed in Denominator** equals No, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Patient encounter during the performance period as listed in Denominator** equals Yes, proceed to check *Encounters conducted via telehealth*.
4. Check *Encounters conducted via telehealth*:
 - a. If *Encounters conducted via telehealth* equals Yes, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Encounters conducted via telehealth* equals No, proceed to check *Place of Service (POS)*.
5. Check *Place of Service (POS)*:
 - a. If *Place of Service (POS)* equals Yes, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Place of Service (POS)* equals No, proceed to check *Documentation stating the patient has received or is currently receiving palliative or hospice care*.
6. Check *Documentation stating the patient has received or is currently receiving palliative or hospice care*:
 - a. If *Documentation stating the patient has received or is currently receiving palliative or hospice care* equals Yes, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Documentation stating the patient has received or is currently receiving palliative or hospice care* equals No, proceed to check *Documentation of patient pregnancy anytime during the measurement period prior to and including the current encounter*.
7. Check *Documentation of patient pregnancy anytime during the measurement period prior to and including the current encounter*:
 - a. If *Documentation of patient pregnancy anytime during the measurement period prior to and including the current encounter* equals Yes, do not include in *Eligible Population/Denominator*. Stop processing.
 - b. If *Documentation of patient pregnancy anytime during the measurement period prior to and including the current encounter* equals No, include in *Eligible Population/Denominator*.

8. Denominator Population

- Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 patients in the Sample Calculation.

9. Start Numerator

10. Check *BMI is documented within normal parameters and no follow-up plan is required*:

- a. If *BMI is documented within normal parameters and no follow-up plan is required* equals Yes, include in *Data Completeness Met and Performance Met*.
 - *Data Completeness Met and Performance Met* letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a¹ equals 30 patients in Sample Calculation.
- b. If *BMI is documented within normal parameters and no follow-up plan is required* equals No, proceed to check *BMI is documented above normal parameters and a follow-up plan is documented*.

11. Check *BMI is documented above normal parameters and a follow-up plan is documented*:

- a. If *BMI is documented above normal parameters and a follow-up plan is documented* equals Yes, include in *Data Completeness Met and Performance Met*.
 - *Data Completeness Met and Performance Met* letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a² equals 20 patients in the Sample Calculation.
- b. If *BMI is documented above normal parameters and a follow-up plan is documented* equals No, proceed to check *BMI is documented below normal parameters and a follow-up plan is documented*.

12. Check *BMI is documented below normal parameters and a follow-up plan is documented*:

- a. If *BMI is documented below normal parameters and a follow-up plan is documented* equals Yes, include in *Data Completeness Met and Performance Met*.
 - *Data Completeness Met and Performance Met* letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a³ equals 0 patients in the Sample Calculation.
- b. If *BMI is documented below normal parameters and a follow-up plan is documented* equals No, proceed to check *BMI not documented due to medical reason OR patient refusal of height or weight measurement*.

13. Check *BMI not documented due to medical reason OR patient refusal of height or weight measurement*:

- a. If *BMI not documented due to medical reason OR patient refusal of height or weight measurement* equals Yes, include in *Data Completeness Met and Denominator Exception*.
 - *Data Completeness Met and Denominator Exception* letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b¹ equals 0 patients in the Sample Calculation.
- b. If *BMI not documented due to medical reason OR patient refusal of height or weight measurement* equals No, proceed to check *BMI is documented as being outside of normal parameters, follow-up plan is not*

completed for documented medical reason.

14. Check *BMI is documented as being outside of normal parameters, follow-up plan is not completed for documented medical reason:*
 - a. If *BMI is documented as being outside of normal parameters, follow-up plan is not completed for documented medical reason* equals Yes, include in *Data Completeness Met and Denominator Exception*.
 - *Data Completeness Met and Denominator Exception* letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b² equals 0 patients in the Sample Calculation.
 - b. If *BMI is documented as being outside of normal parameters, follow-up plan is not completed for documented medical reason* equals No, proceed to check *BMI not documented and no reason is given*.
15. Check *BMI not documented and no reason is given:*
 - a. If *BMI not documented and no reason is given* equals Yes, include in *Data Completeness Met and Performance Not Met*
 - *Data Completeness Met and Performance Not Met* letter is represented as Data Completeness in the Sample Calculation listed at the end of this document. Letter c¹ equals 10 patients in the Sample Calculation.
 - b. If *BMI not documented and no reason is given* equals No, proceed to check *BMI documented outside of normal parameters, no follow-up plan documented, no reason given*.
16. Check *BMI documented outside normal parameters, no follow-up plan documented, no reason given:*
 - a. If *BMI documented outside normal parameters, no follow-up plan documented, no reason given* equals Yes, include in *Data Completeness Met and Performance Not Met*.
 - *Data Completeness Met and Performance Not Met* letter is represented as Data Completeness in the Sample Calculation listed at the end of this document. Letter c² equals 10 patients in the Sample Calculation.
 - b. If *BMI documented outside normal parameters, no follow-up plan documented, no reason given* equals No, proceed to check *Data Completeness Not Met*.
17. Check *Data Completeness Not Met*:
 - If *Data Completeness Not Met*, Quality Data Code or equivalent not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

Sample Calculations:

Data Completeness equals Performance Met (a¹ plus a² plus a³ equals 50 patients) plus Denominator Exception (b¹ plus b² equals 0 patients) plus Performance Not Met (c¹ plus c² equals 20 patients) divided by Eligible Population / Denominator (d equals 80 patients). All equals 70 patients divided by 80 patients. All equals 87.50 percent.

Performance Rate equals Performance Met (a¹ plus a² plus a³ equals 50 patients) divided by Data Completeness Numerator (70 patients) minus Denominator Exception (b¹ plus b² equals 0 patients). All equals 50 patients divided by 70 patients. All equals 71.43 percent.

* See the posted measure specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Patient-Intermediate

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.